

Being Fat and Friendly is Not Enough

Sariel Har-Peled*

August 17, 2009

Abstract

We show that there is no $(1 + \varepsilon)$ -approximation algorithm for the problem of covering points in the plane by minimum number of fat triangles of similar size (with the minimum angle of the triangles being close to 45 degrees). Here, the available triangles are prespecified in advance. Since a constant factor approximation algorithm is known for this problem [CV07], this settles the approximability of this problem.

We also investigate some related problems, including cover by friendly fat shapes, and independent set of triangles in three dimensions.

1 Introduction

In the *planar set-cover* problem, we are given a set Q of points in the plane, and a set of shapes S , and we would like to find a minimum cardinality subset of shapes of S , such that their union covers all the points of Q .

In Section 2, we show that this problem is **MAXSNP-HARD** even when the shapes are fat, convex, of the same size, and their union complexity is linear. As such, no PTAS for such a problem is possible (unless $P = NP$). We remind the reader that a PTAS (Polynomial Time Approximation Scheme) is an approximation algorithm that given an input and a parameter $\varepsilon > 0$ it outputs a $(1 + \varepsilon)$ -approximate solution to the given instance in polynomial time.

In fact, the same result holds even if the shapes are fat triangles, with minimum angle approaching 45 degrees and of similar size. In fact, in this case, all the triangles are rotated and translated “noisy” copies of the same triangle, where the original triangle is right-angled and also isosceles, with the two base angles being 45 degrees. (The word “noisy” here means that we can make all the lengths of the edges of the copies of this triangle to be arbitrarily close to the lengths of the corresponding original edges.)

Finally, in Section 4, we show that there is no PTAS for independent set of triangles in $\mathbb{R}^3$. Here, we are given a set of triangles in three dimensions, and we are looking for the largest subset of triangles such that no pair of them intersects.

*Department of Computer Science; University of Illinois; 201 N. Goodwin Avenue; Urbana, IL, 61801, USA; sariel@uiuc.edu; <http://www.uiuc.edu/~sariel/>.

Known results. For relevant results see [AES09, CCH09, MR09] and references therein. In particular, our basic construction of Section 2, is similar to one of the hardness proofs of [EvL08]. Several **APX-HARDNESS** results are known in geometry, among them hardness of (i) independent set of boxes in 3d [CC05], (ii) maximizing guarded boundary in art gallery problems [FMZ07], and (iii) separating points by axis-parallel lines [CDKW05]. This list is by no means exhaustive.

2 Hardness of approximation of the friendly geometric set-cover problem

Consider a set system $(U, \mathcal{F})$, where $|U| = n$, and every element of $\mathcal{F}$ is a subset of U of size at most k , where k is some fixed constant. We are interested in finding the minimum cardinality cover of U by sets of $\mathcal{F}$. This is known as the **minimum k -set cover** problem and it is **MAXSNP-HARD** for $k \geq 3$ [ACG⁺99]. It is known that if a problem is **MAXSNP-HARD** then there is no PTAS for it unless $\mathbf{P} = \mathbf{NP}$. Here, one can even assume that every point of U participates in at most $k + 1$ sets of $\mathcal{F}$.

Definition 2.1 Let $\mathbf{Q}$ be a set of n points in the plane, and let $\mathcal{F}$ be a set of m regions in the plane, such that

- (i) the shapes of $\mathcal{F}$ are convex, fat, and of similar size,
- (ii) the boundaries of any pair of shapes of $\mathcal{F}$ intersect in at most 6 points,
- (iii) the union complexity of any m shapes of $\mathcal{F}$ is $O(m)$,
- (iv) and, any point of $\mathbf{Q}$ is covered by a constant number of shapes of $\mathcal{F}$.

We are interested in the problem of finding a minimum number of shapes of $\mathcal{F}$ that covers all the points of $\mathbf{Q}$. We will refer to this variant as the **friendly geometric set cover** problem.

Lemma 2.2 *There is no PTAS for the friendly geometric set cover problem, unless $\mathbf{P} = \mathbf{NP}$.*

Proof: We will reduce an instance $(U, \mathcal{F})$ of the minimum k -set cover problem (for $k = 3$) into an instance of the friendly geometric set cover problem. So, let $U = \{u_1, \dots, u_n\}$, and $\mathcal{F} = \{S_1, \dots, S_m\}$. We place n points equally spaced on the unit radius circle centered at the origin, and let $\mathbf{Q} = \{\mathbf{p}_1, \dots, \mathbf{p}_n\}$ be the resulting set of points. Let $f(u_i) = \mathbf{p}_i$, for $i = 1, \dots, n$. Next, we map the set S_i (which is of size at most k) to the region

$$R_i = \mathcal{CH} \left(\text{disk} \left(1 - \frac{i}{10n^2m} \right) \cup f(S_i) \right),$$

for $i = 1, \dots, m$, where $\mathcal{CH}$ is the convex hull, $f(S_i) = \cup_{x \in S_i} \{f(x)\}$, and $\text{disk}(r)$ denotes the disk of radius r centered at the origin. Visually, R_i is a disk with three (since $k = 3$) teeth coming out of it, see Figure 1. Note, that the boundary of two such shapes intersects in at most 6 points.




Figure 1: (i) A region R constructed for the set $S_t = \{u_i, u_j, u_k\}$. Observe that in the construction, the inner disk is even bigger. As such, no two points are connected by an edge of the convex-hull when we add in the inner disk to the convex-hull. As such, each point “contribution” to the region R is separated from the contribution of other points. (ii) How the intersection of the boundary of two such regions looks like.

It is now easy to verify that the resulting instance of geometric set cover $(Q, \{R_1, \dots, R_m\})$ is in fact friendly, and clearly any cover of Q by these shapes can be interpreted as a cover of U by the corresponding sets of $\mathcal{F}$. Thus, a PTAS for the friendly geometric set cover problem, would imply a PTAS for the minimum k -set cover, which is impossible unless $P = NP$. ■

3 Hardness of approximation for set-cover by fat triangles

It is known that **Vertex Cover** is **APX-HARD** even for a graph with a maximum degree 3 [ACG⁺99]. A problem that is **APX-HARD** does not have a PTAS unless $P = NP$. Consider such a graph G , and observe that a **Vertex Cover** problem in such a graph can be reduced to **Set Cover** where every set is of size at most 3. Indeed, the ground set U is the edges of G , and every vertex $v \in V(G)$ gives a rise to the set $S_v = \{e \mid v \in e \text{ and } e \in E(G)\}$, which is of size at most 3. Clearly, any cover C of size t for the set system $\mathcal{X} = (U, \{S_v \mid v \in V(G)\})$, has a corresponding vertex cover of G of the same size. Thus, **Set Cover** with every set of size (at most) three is **APX-HARD** (this is of course well known). Note, that in this set cover instance, every element participate in exactly two sets (i.e., the two vertices adjacent to the original edge).

The graph G is of maximum degree three, and by Vizing’s theorem [BM76], it is 4 edge-colorable¹. Thus, the ground set of the set system $\mathcal{X}$ can be colored by 4 colors, and no set in this set system has a color appearing more than once.

In the **fat-triangle set cover problem**, specified by a set of points in the plane Q and a set of fat triangles $\mathcal{T}$, one wants to find the minimum subset of $\mathcal{T}$ such that its union covers

¹Vizing’s theorem states that a graph with maximum degree Δ can be edge colored by $\Delta + 1$ colors. In this specific case, one can reach the same conclusion directly from Brook’s theorem.

all the points of $\mathcal{Q}$.

Lemma 3.1 *There is no PTAS for the fat-triangle set cover problem, unless $\mathsf{P} = \mathsf{NP}$.*

In fact, one can prespecify an arbitrary constant $\delta > 0$, and the claim would hold true even if the following conditions hold on the given instance $(\mathcal{Q}, \mathcal{T})$:

- (A) *The minimum angle of all the triangles of $\mathcal{T}$ is larger than $45 - \delta$ degrees.*
- (B) *No point of $\mathcal{Q}$ is covered by more than two triangles of $\mathcal{T}$.*
- (C) *The points of $\mathcal{Q}$ are in convex position.*
- (D) *All the triangles of $\mathcal{T}$ are of similar size. Specifically, all the triangles diameter is in the range (say) $(2 - \delta, 2]$.*
- (E) *Each triangle of $\mathcal{T}$ has two angles in the range $(45 - \delta, 45 + \delta)$, and one angle in the range $(90 - \delta, 90 + \delta)$.*
- (F) *The vertices of the triangles of $\mathcal{T}$ are the points of $\mathcal{Q}$.*


Proof: We are given an instance of the vertex cover problem for a graph with maximum degree 3, and we transform it into a set cover instance as mentioned above, denoted by $\mathcal{X} = (U, \mathcal{F}_{\mathcal{X}})$. Let $n = |U|$, and color U (as described above) by 4 colors such that no set of $\mathcal{X}$ has the same color repeated twice, let $U_1, \dots, U_4$ be the partition of U by the color of the points.

Let $\mathcal{C}$ denote the circle of radius one centered at the origin. We place four relatively short arcs on $\mathcal{C}$, placed on the four intersection points of $\mathcal{C}$ with the x and y axes, see figure on the right. Let $I_1, \dots, I_4$ denote these four circular intervals. We equally space the elements of U_i (as points) on the interval I_i , for $i = 1, \dots, 4$. Let $\mathcal{Q}$ be the resulting set of points.

For every set $S \in \mathcal{F}_{\mathcal{X}}$, take the convex hull of the points corresponding to its elements as its representing triangle T_S . Note, that since the vertices of T_S lie on three intervals out of I_1, I_2, I_3, I_4 , it follows that it must be fat, for all $S \in \mathcal{F}_{\mathcal{X}}$. As such, the resulting set of triangles $\mathcal{T} = \{T_S \mid S \in \mathcal{F}_{\mathcal{X}}\}$ is fat, and clearly there is a cover of $\mathcal{Q}$ by t triangles of $\mathcal{T}$ if and only if the original set cover problem has a cover of size t .

In fact, any triangle having its three vertices on three different intervals of $I_1, \dots, I_4$ is close to being an isosceles triangle with the middle angle being 90 degrees. As such, by choosing these intervals to be sufficiently short, any triangle of $\mathcal{T}$ would have a minimum degree larger than, say, $45 - \delta$ degrees, and with diameter in the range between $2 - \delta$ and 2.

This is clearly an instance the fat-triangle set cover problem. Solving it is equivalent to solving the original vertex cover problem, but since it is $\mathsf{APX-HARD}$, it follows that the fat-triangle set cover problem is $\mathsf{APX-HARD}$. $\blacksquare$



Remark 3.2 For fat triangles of similar size a constant factor approximation algorithm is known [CV07]. Lemma 3.1 implies that one can do no better. Naturally, it might be possible to slightly improve the constant of approximation provided by the algorithm of Clarkson and Varadarajan [CV07].

However, for fat triangles of different sizes, only a $\log \log \log$ approximation is known [AES09]. It is natural to ask if this can be improved.

3.1 Extensions

Lemma 3.3 *Given a set of points Q in the plane and a set of circles $\mathcal{F}$, finding the minimum number of circles of $\mathcal{F}$ that covers Q is **APX-HARD**; that is, there is no PTAS for this problem.*

Proof: Slightly perturb the point set used in the proof of Lemma 3.1, so that no four points of it are co-circular. Let Q denote the resulting set of points. For every set $S \in \mathcal{F}_X$, we now take the circle passing through the three corresponding points. Clearly, this results in a set of circles (that are almost identical, but yet all different), such that finding the minimum number of circles covering the set Q is equivalent to solving the original problem. ■

Lemma 3.4 *Given a set of points P in $\mathbb{R}^3$ and a set of planes $\mathcal{F}$, finding the minimum number of planes of $\mathcal{F}$ that covers P is **APX-HARD**; that is, there is no PTAS for this problem.*

Proof: Let Q be the point set and $\mathcal{F}$ be the set of circles constructed in the proof of Lemma 3.3, and map every point in it to three dimensions using the mapping $f : (x, y) \rightarrow (x, y, x^2 + y^2)$. This is a standard lifting map used in computing planar Delaunay triangulations via convex-hull in three dimensions, see [dBvKOS00]. Let $P = f(Q)$ be the resulting point set.

It is easy to verify that a circle of $c \in \mathcal{F}$ is mapped by f into a curve that lies on a plane. We will abuse notations slightly, and use $f(c)$ to denote this plane. Let $\mathcal{H} = f(\mathcal{F})$. Furthermore, for a circle $c \in \mathcal{F}$, we have that $f(c \cap Q) = f(c) \cap P$. Namely, solving the set cover problem $(P, \mathcal{H})$ is equivalent to solving the original set cover instance $(Q, \mathcal{F})$. ■

Interestingly, the recent work of Mustafa and Ray [MR09] implies that there is a PTAS for set cover of points by disks (i.e., circles with their interior), and similarly, there is a PTAS for the problem of set cover of points by half-spaces in three dimensions. Thus, somewhat surprisingly, the “shelled” version of these problems are harder than the filled-in version.

4 Hardness of independent set of triangles in 3d

Given a set S of n objects in $\mathbb{R}^d$ (say, triangles in 3d), we are interested in computing a maximum number of objects that are **independent**; that is, no pair of objects in this set (i.e., independent set) intersects. This is the geometric realization of the **independent set** problem for the intersection graph induced by these objects.

Lemma 4.1 *There is no PTAS for the maximum independent set of triangles in $\mathbb{R}^3$, unless $P = NP$.*

Proof: Independent set is **APX-HARD** even for graphs with maximum degree 3 [ACG⁺99]. So, let $G = (V, E)$ be a given such graph with maximum degree 3, where $V = \{v_1, \dots, v_n\}$. We will create a set of triangles, such that their intersection graph is G .

We use the following fact: If one spreads n points $p_1, \dots, p_n$ on the positive branch of the moment curve in $\mathbb{R}^3$ [Sei91, EK03], their Voronoi diagram is **neighborly**; that is, every Voronoi cell is a convex polytope that shares a non-empty two dimensional boundary face with each of the other cells of the diagram. Let C_i denote the cell of the point p_i in this Voronoi diagram, for $i = 1, \dots, n$.

Now, for every vertex $v_i \in V$, we form a set Q_i of (at most) three points, as follows. If $v_i v_j \in E$, then we place a point p_{ij} on the common boundary of C_i and C_j , and we add this point to Q_i and Q_j .

For $i = 1, \dots, n$, the region f_i corresponding to v_i is the triangle formed by the convex-hull of Q_i (if Q_i have fewer than three points then the triangle is degenerate).

Let $\mathcal{T} = \{f_1, \dots, f_n\}$. Observe that the triangles of $\mathcal{T}$ are disjoint except maybe in their common vertices, as their interior is contained inside the interior of C_i , and the cells $C_1, \dots, C_n$ are interior disjoint. Clearly $f_i \cap f_j \neq \emptyset$ if and only if $v_i v_j \in E$. Thus, finding an independent set in G is equivalent to finding an independent set of triangles of the same size in $\mathcal{T}$. We conclude that the problem of finding maximum independent set of triangles is **APX-HARD**, and as such does not have a PTAS unless $P = NP$. ■

The above result is still a far cry from being tight. In light of the results known [AM06] for independent set of segments in the plane, it is natural to conjecture that this harder problem (i.e., finding independent set of triangle) can not be approximated to within any polynomial factor.

Implicit in the above proof is the fact that any graph can be realized as the intersection graph of convex bodies in $\mathbb{R}^3$ (we were a bit more elaborate for the sake of completeness and since we needed slightly more structure). This is well known and can be traced to a result of Tietze from 1905 [Tie05].

Acknowledgments

The author thanks Janos Pach and Shakhar Smorodinsky for useful discussions on this problem.

References

- [ACG⁺99] G. Ausiello, P. Crescenzi, G. Gambosi, V. Kann, A. Marchetti-Spaccamela, and M. Protasi. *Complexity and approximation*. Springer-Verlag, Berlin, 1999.
- [AES09] B. Aronov, E. Ezra, and M. Sharir. Small-size eps-nets for axis-parallel rectangles and boxes. In *Proc. 41th Annu. ACM Sympos. Theory Comput.*, 2009.

- [AM06] P. K. Agarwal and N. H. Mustafa. Independent set of intersection graphs of convex objects in 2D. *Comput. Geom. Theory Appl.*, 34(2):83–95, 2006.
- [BM76] J. A. Bondy and U. S. R. Murty. *Graph Theory with Applications*. North-Holland, 1976.
- [CC05] M. Chlebík and J. Chlebíková. Approximation hardness of optimization problems in intersection graphs of d -dimensional boxes. In *Proc. 16th ACM-SIAM Sympos. Discrete Algorithms*, pages 267–276, 2005.
- [CCH09] C. Chekuri, K. L. Clarkson., and S. Har-Peled. On the set multi-cover problem in geometric settings. In *Proc. 25th Annu. ACM Sympos. Comput. Geom.*, pages 341–350, 2009.
- [CDKW05] G. Calinescu, A. Dumitrescu, H. J. Karloff, and P. Wan. Separating points by axis-parallel lines. *Internat. J. Comput. Geom. Appl.*, 15(6):575–590, 2005.
- [CV07] K. L. Clarkson and K. R. Varadarajan. Improved approximation algorithms for geometric set cover. *Discrete Comput. Geom.*, 37(1):43–58, 2007.
- [dBvKOS00] M. de Berg, M. van Kreveld, M. H. Overmars, and O. Schwarzkopf. *Computational Geometry: Algorithms and Applications*. Springer-Verlag, 2nd edition, 2000.
- [EK03] J. Erickson and S. Kim. Arbitrarily large neighborly families of congruent symmetric convex 3-polytopes. In A. Bezdek, editor, *Discrete Geometry: In Honor of W. Kuperberg’s 60th Birthday*, Lecture Notes Pure Appl. Math., pages 267–278. Marcel-Dekker, 2003.
- [EvL08] T. Erlebach and E. J. van Leeuwen. Domination in geometric intersection graphs. In *8th Latin Amer. Theo. Inf.*, pages 747–758, 2008.
- [FMZ07] C. Fragoudakis, E. Markou, and S. Zachos. Maximizing the guarded boundary of an art gallery is apx-complete. *Comput. Geom. Theory Appl.*, 38(3):170–180, 2007.
- [MR09] N. H. Mustafa and S. Ray. Improved results on geometric hitting set problems. In *Proc. 25th Annu. ACM Sympos. Comput. Geom.*, 2009. to appear.
- [Sei91] R. Seidel. Exact upper bounds for the number of faces in d -dimensional Voronoi diagrams. In P. Gritzman and B. Sturmfels, editors, *Applied Geometry and Discrete Mathematics: The Victor Klee Festschrift*, volume 4 of *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, pages 517–530. AMS Press, 1991.
- [Tie05] H. Tietze. Über das problem der nachbargebiete im raum. *Monatshefte Math.*, 10:211–216, 1905.